

Weatherman-ager Managing Weather Risk

Oxbow, Saskatchewan

Management of fertility, crop residue, weeds, insects and diseases are all part of good agronomic practice in Prairie Agriculture. However, producers have traditionally reacted to climatic variability rather than managing it. Many producers manage the economic influence of climatic variability through participation in crop and hail insurance, but to date the only agronomic management tools have been in the form of early maturing crops.

The *Weatherman-ager* is a management tool that gives you access to 49 years of weather records. It has been developed to predict the probability of crops reaching maturity based on different seeding dates.

What is physiological maturity?

Physiological maturity is the point at which maximum kernel dry weight is achieved. Physiological maturity coincides with the earliest time a crop could be sprayed with pre-harvest glyphosate or swathed.

Actual harvest date typically occurs 10 to 14 days after physiological maturity date.

How does a probability table work?

The probability tables included in this document depict alternative seeding dates on one axis and physiological maturity date on the other axis.

The values within the table indicate how many years out of 100 a crop would reach physiological maturity for that seeding date.

How were the probability tables developed?

Weather data from Environment Canada was analysed from 1949 to 1998 for selected weather stations. Growing Degree Day (GDD) accumulations were then calculated for each station based on a variety of seeding dates.

What is a Growing Degree Day and how is it calculated?

Growing Degree Days are a commonly used method of measuring and displaying heat accumulation and crop heat requirements. An example of the calculation utilized to measure growing degree accumulation for a summer day with a high of 20 Celsius and a low of 10 Celsius would be as follows:

$$\begin{aligned} \text{GDD} &= (20(\text{daily high}) + 10(\text{daily low}) / 2) - 5 \\ &= (30 / 2) - 5 \\ &= 15 - 5 \\ &= 10 \text{ GDD accumulated during our example summer day.} \end{aligned}$$

- * All temperatures are in degrees Celsius.
- * Five is subtracted from the average of the daily high and low temperatures, as 5 Celsius is the generally accepted minimum temperature requirement for plant growth and development.

If you have comments or questions please contact a Ducks Unlimited Canada Agriologist at (306)359-2229 or Winter Cereals Canada at info@wintercereals.com.



What are GDD requirements for crops grown in Western Canada?

Research done by Dr. Brian Fowler and Brian Duggan at the University of Saskatchewan has shown little geographic variability in GDD requirements for crops throughout Western Canada. For example, Argentine Canola grown in Prince Albert and Shaunavon require approximately the same 1040 GDD to reach physiological maturity. Physiological maturity is determined primarily by heat accumulation; moisture levels only affect physiological maturity under extreme drought conditions.

Crop	GDD Requirement
Flax	1200
HRSW	1175
Argentine Canola	1040
Mustard	1004
Oats	961
Barley	850
Polish Canola	850

GDD requirements shown are for an average of varieties commonly grown in Prairie Canada.

An example of using the *Weatherman-ager* to plan winter wheat seeding would be:

Recommended seeding window for Winter Wheat seeding is Aug. 25 to Sept. 15.

Optimum date for seeding Winter Wheat at Oxbow, SK: September 6

To have stubble available following a crop of Argentine Canola:

Estimated harvest date: Aug. 30 to Sept. 3

Probability of physiological maturity by Aug. 20: 80%

Argentine Canola seeding date to achieve winter wheat seeding date: May 20

This table can also be utilized to determine the probability of achieving physiological maturity prior to first fall frost.

An example of using the *Weatherman-ager* to plan spring seeding would be:

Average first fall frost at Estevan, SK. (Closest available recording) – September 21

Argentine Canola seeding date necessary to avoid average first frost date: June 9

Example: Argentine Canola at Oxbow, Saskatchewan

		Physiological Maturity Date									
		05-Aug	10-Aug	15-Aug	20-Aug	25-Aug	30-Aug	04-Sep	09-Sep	14-Sep	19-Sep
Seed Date	01-May	32	64	84	94	98	98	98	98	98	98
	05-May	28	62	78	86	98	98	98	98	98	98
	10-May	22	56	74	82	98	98	98	98	98	98
	15-May	14	44	68	82	92	98	98	98	98	98
	20-May	6	24	54	80	84	96	98	98	98	98
	25-May	4	16	32	58	82	92	94	98	98	98
	30-May	2	4	18	50	70	82	92	98	98	98
	04-Jun	0	2	12	22	50	76	84	88	98	98
	09-Jun	0	0	2	12	28	52	74	82	88	94

The *Weatherman-ager* tables were generated using Environment Canada weather station data collected from 1949 to 1998.

If you have comments or questions please contact a Ducks Unlimited Canada Agrologist at (306)359-2229 or Winter Cereals Canada at info@wintercereals.com.



Oxbow, Saskatchewan (4015800)

The table below was generated from Environment Canada weather station data collected from 1949 to 1998.

Crops:

Argentine Canola

Physiological Maturity Date

	05-Aug	10-Aug	15-Aug	20-Aug	25-Aug	30-Aug	04-Sep	09-Sep	14-Sep	19-Sep
01-May	32	64	84	94	98	98	98	98	98	98
05-May	28	62	78	86	98	98	98	98	98	98
10-May	22	56	74	82	98	98	98	98	98	98
15-May	14	44	68	82	92	98	98	98	98	98
20-May	6	24	54	80	84	96	98	98	98	98
25-May	4	16	32	58	82	92	94	98	98	98
30-May	2	4	18	50	70	82	92	98	98	98
04-Jun	0	2	12	22	50	76	84	88	98	98
09-Jun	0	0	2	12	28	52	74	82	88	94

Flax

Physiological Maturity Date

	05-Aug	10-Aug	15-Aug	20-Aug	25-Aug	30-Aug	04-Sep	09-Sep	14-Sep	19-Sep
01-May	4	10	24	54	74	88	92	96	98	98
05-May	2	10	18	44	72	80	88	96	98	98
10-May	2	4	16	36	66	78	82	92	98	98
15-May	2	2	12	26	52	76	78	84	94	96
20-May	0	2	8	16	36	56	76	80	86	94
25-May	0	2	2	12	22	42	66	76	82	84
30-May	0	0	2	2	16	28	44	64	72	82
04-Jun	0	0	0	2	10	18	28	46	62	70
09-Jun	0	0	0	0	0	10	20	30	42	56

HRS Wheat

Physiological Maturity Date

	05-Aug	10-Aug	15-Aug	20-Aug	25-Aug	30-Aug	04-Sep	09-Sep	14-Sep	19-Sep
01-May	6	14	36	62	80	90	98	98	98	98
05-May	4	10	32	56	76	84	96	96	98	98
10-May	2	8	20	46	72	82	92	96	98	98
15-May	2	4	14	32	62	78	82	92	96	98
20-May	2	2	10	20	50	70	80	86	92	96
25-May	0	2	4	14	32	54	74	80	84	88
30-May	0	0	2	12	18	36	58	68	78	84
04-Jun	0	0	0	2	12	22	36	56	66	78
09-Jun	0	0	0	0	2	12	24	36	58	60

If you have comments or questions please contact a Ducks Unlimited Canada Agrolgist at (306)359-2229 or Winter Cereals Canada at info@wintercereals.com.

Mustard

Physiological Maturity Date

	05-Aug	10-Aug	15-Aug	20-Aug	25-Aug	30-Aug	04-Sep	09-Sep	14-Sep	19-Sep
01-May	56	76	90	96	98	98	98	98	98	98
05-May	42	68	84	96	98	98	98	98	98	98
10-May	36	64	78	92	98	98	98	98	98	98
15-May	22	60	76	84	98	98	98	98	98	98
20-May	14	42	68	82	96	98	98	98	98	98
25-May	4	22	58	80	86	94	98	98	98	98
30-May	4	16	30	60	80	90	96	98	98	98
04-Jun	0	4	14	42	70	86	92	98	98	98
09-Jun	0	0	6	18	46	74	82	88	98	98

Oat

Physiological Maturity Date

	05-Aug	10-Aug	15-Aug	20-Aug	25-Aug	30-Aug	04-Sep	09-Sep	14-Sep	19-Sep
01-May	68	90	96	98	98	98	98	98	98	98
05-May	62	86	96	98	98	98	98	98	98	98
10-May	58	74	90	98	98	98	98	98	98	98
15-May	50	68	86	96	98	98	98	98	98	98
20-May	30	64	82	92	98	98	98	98	98	98
25-May	14	46	66	82	94	98	98	98	98	98
30-May	6	22	56	78	88	96	98	98	98	98
04-Jun	2	14	30	64	76	92	98	98	98	98
09-Jun	0	0	12	40	68	82	90	98	98	98

Polish canola / Barley

Physiological Maturity Date

	05-Aug	10-Aug	15-Aug	20-Aug	25-Aug	30-Aug	04-Sep	09-Sep	14-Sep	19-Sep
01-May	96	98	98	98	98	98	98	98	98	98
05-May	92	98	98	98	98	98	98	98	98	98
10-May	90	96	98	98	98	98	98	98	98	98
15-May	84	94	98	98	98	98	98	98	98	98
20-May	72	90	98	98	98	98	98	98	98	98
25-May	62	84	94	98	98	98	98	98	98	98
30-May	42	74	86	98	98	98	98	98	98	98
04-Jun	20	54	78	94	98	98	98	98	98	98
09-Jun	8	34	62	78	98	98	98	98	98	98

If you have comments or questions please contact a Ducks Unlimited Canada Agriologist at (306)359-2229 or Winter Cereals Canada at info@wintercereals.com.